

M5 Anti-HA-Tag mAb (Agarose conjugated)使用说明书

产品名称	单位	货号	Beads 含量
M5 Anti-HA-Tag mAb (Agarose conjugated)	1ml	MF097-01	0.5ml
M5 Anti-HA-Tag mAb (Agarose conjugated)	5ml	MF097-05	2.5ml

【STORAGE】

The product is supplied as a 50% slurry in storage buffer (1 PBS, pH 7.4, containing 0.1% NaN₃).

Store the product at 4°C and do not freeze.

【BACKGROUND】

Anti-HA-Tag Mouse mAb (Agarose Conjugated) is a monoclonal anti-HA antibody covalently linked to agarose; the agarose enables immunoprecipitation (IP) of HA tagged proteins or co-immunoprecipitation (Co-IP) of their interacting partners.

【SOURCE】


This monoclonal antibody is produced by immunizing animals with a synthetic peptide containing the influenza hemagglutinin epitope (YPYDVPDYA) (KLH-coupled).

【SPECIFICITY】 : Anti-HA-Tag Mouse mAb detects transfected proteins containing the HA epitope tag.

【REACTIVITY】: All

【ISOTYPE】 : Mouse IgG1

【RECOMMENDED ELUTION BUFFER】 : 0.2 M Glycine, pH 2.2



HEK 293T cells were transfected with HA-tagged protein or not, and 100 μ l cell lysate (about 100 μ g total protein) was incubated with 30 μ l 50% slurry of Anti-HA Agarose for 3 h at 4°C. After washing, the beads were eluted by 30 μ l elution buffer twice. After neutralization of the eluant, 6 μ l 6 \times SDS loading buffer was added. Then 20 μ l sample was subjected to the SDS-PAGE. Blot was probed with Anti-HA-Tag Mouse mAb.

Lane 1: 1st Elution with elution buffer.

Lane 2: IP of untransfected HEK 293T lysate.

【IMMUNOPRECIPITATION PROCEDURE】

The work can be performed in 1.5 ml micro-centrifuge tubes or in spin columns.

1. Thoroughly resuspend the Anti-HA Agarose by inverting the tube or vial several times.
2. Add 20-50 μ l 50% slurry of Anti-HA Agarose into cell lysate using a widebore pipette tip.

Note: The lysate should be fresh, and for a well expressed tagged protein, 200 μ l lysate (200-500 μ g total protein) usually yields a good IP result.

3. Incubate with gentle mixing for 2 h to overnight at 4°C.
4. Wash the beads with 1 ml TBS buffer or lysis buffer, such as RIPA (50 mM Tris HCl, pH 7.4, 150 mM NaCl, 1 mM EDTA, 1% NP-40, 0.5% sodium deoxycholate), centrifuge for 3 min at 2,000x g, and discard the supernatant. Wash 3 times, avoid losing beads during washes.
5. Elution of the HA tagged protein.

Option 1. Elution with elution buffer.

Add 30-50 μ l elution buffer to the beads, gently tap the tube to mix well, immediately centrifuge for 3 min, transfer the supernatant very carefully to a fresh tube (Avoid transferring any beads).

Note: Neutralize the eluant immediately by add 1 μ l of 1.5 M Tris, pH 9.0 per 20 μ l Elution buffer.

Option 2. Elution with HA peptide

Add 30-50 μ l HA peptide solution (100 μ g/ml HA peptide in TBS buffer), gently tap the tube to mix well, incubate for 10 min, centrifuge for 3 min, and transfer the supernatant to a fresh tube. TBS buffer: 50 mM Tris HCl, 150 mM NaCl, pH 7.4.

Option 3. Elution with SDS loading buffer

Add 30 μ l 2x SDS loading buffer, gently tap the tube to mix well, boil at 100°C for 5 min, centrifuge for 3 min, transfer the supernatant to a fresh tube.

Note: in this case, the supernatant contains not only the binding proteins, but also IgG (heavy and light chains).

6. Prepare SDS-PAGE gel for western blotting or proceed to other assays.

【备注】

本产品仅供科研使用。在确认产品质量出现问题时，本公司承诺为客户免费更换等量的质量合格产品。